

Atascadero Pet Hospital and Emergency Center
9575 El Camino Real Atascadero, CA 93422
(805) 466-3880 (805) 466-3830 Fax
www.apetcenter.com

FELINE TAPEWORM INFECTION

Tapeworms are intestinal parasites of the cat and dog. Because they are classified as *cestodes*, they belong to a different family than the hookworms and roundworms which are called *nematodes*. Several types of tapeworms are known to infect cats. *Dipylidium caninum* is by far the most common one.

The tapeworm uses its hook-like mouthparts for anchoring to the wall of the small intestine. Eventually, adult tapeworms may reach several inches in length. As the adult matures, individual segments called *proglottids* break off from the main body of the tapeworm and pass into the cat's feces.


Are certain cats more likely to get tapeworms?

Fleas are the intermediate host for the *Dipylidium caninum*. In other words, the tapeworm is unable to complete its life cycle without the presence of fleas in the environment. Regardless of whether the owner may have seen fleas, the cat *must* have ingested a flea in order to have tapeworms. Consequently, tapeworms are more common in environments that are heavily infested with fleas. Lice are also reported as intermediate hosts for tapeworms but they are relatively uncommon parasites of cats.

How do cats get tapeworms?

First, tapeworm eggs must be ingested by flea larvae, an immature stage of the flea. Contact between flea larvae and tapeworm eggs is facilitated by contaminated bedding or carpet. Adult fleas do not participate in this part of the tapeworm lifecycle. The infected flea larvae will mature into adult fleas.

Next, the cat chews or licks its skin as an adult flea bites and swallows the flea. As the flea is digested within the cat's intestine, the tapeworm hatches and anchors itself to the intestinal lining.



What are the clinical signs of tapeworm infection?

Tapeworms are not particularly harmful to the cat and few clinical signs are attributed to their presence. Usually, the cat is presented because of the owner's aversion to the presence of the crawling proglottids. Rarely, tapeworms may cause debilitation or weight loss if they are present in large numbers. A cat will occasionally scoot or drag its anus across the ground or carpet due to the anal irritation caused by the proglottids. However, this behavior is much more common in dogs than cats.

Occasionally, a tapeworm will release its attachment in the intestines and migrate to the stomach. When this happens, the cat may vomit an adult tapeworm several inches in length.


How are tapeworms diagnosed?

Most commonly, owners recognize that the cat has tapeworms and bring this to the attention of the veterinarian. When terminal segments of the tapeworm break off and pass into the cat's stool, they can be seen crawling on the surface of the feces. These proglottid segments look like grains of cooked white rice or cucumber seeds. Less commonly, they are seen crawling around the cat's anus. Each of these proglottid capsules contains up to twenty tapeworm eggs.

When these proglottids are released into the environment, they dehydrate and harden, becoming smaller and taking on a golden hue. Eventually, they break open and release their contents. Be aware that tapeworms are not readily diagnosed with routine fecal examinations. Because of this, you should notify your veterinarian when tapeworm segments are found in your cat's stool.

What is the treatment for tapeworms?

Tapeworm treatments are safe and effective. The deworming medication called an *anthelmintic* may be given as a tablet or an injection. After treatment, the tapeworm dies and is usually digested within the intestine, so worm segments don't usually pass into the stool. Side-effects, such as vomiting and diarrhea, are rarely reported with the newer tapeworm medications.



Flea control is the cornerstone of preventing tapeworm infection. With the new and exciting flea control products that have become available, this is now much easier than ever before. Depending on the type of product you use and the presence of other pets in your home, you may also need to treat your house and yard for fleas. With some of the newer products, environmental control of fleas may not be needed. Circumstances vary, however, so be sure to talk to your veterinarian.

If the cat lives in a flea-infested environment, tapeworm infection can become re-established within a few weeks. This is very rarely due to treatment failure; in most cases, reappearance of tapeworms represents reinfection of the cat. Additional recommendations include:

1. Prompt treatment when tapeworms are detected.

2. Appropriate disposal of all pet feces, especially in public parks, yard, or playgrounds.
3. Strict hygiene practices for children after playing outdoors.

What is the prognosis?

New medications are safe and effective. The prognosis for successful treatment is excellent. Prevention is successfully accomplished by using a monthly flea preventive

Are cat tapeworms a danger to me or my family?

Humans can become infected with tapeworms, although infection is rare because it requires ingestion of a flea. Most reported cases have involved children. The risk for human infection with *Dipylidium caninum* is quite small but does exist.

What are the other tapeworms that can infect my cat?

The other common tapeworms that can infect a cat are members of a group called *Taenia*. The intermediate host of these tapeworms is mice, birds, or rabbits. In a similar manner to *Dipylidium* transmission, cats acquire *Taenia* infestations by eating infected mice, birds, or rabbits.

Another less common group of tapeworms called *Echinococcus* is of increasing concern as a threat to human health. These tapeworms cause serious, potentially fatal, disease when humans become infected. Infection with this parasite is harder to diagnose than *Dipylidium* because the segments are small and not readily seen. Trappers and hunters in the north central United States and south central Canada may be at increased risk for infection with this worm when strict hygiene is not practiced. Foxes and coyotes and the wild rodents upon which they prey are important in the life cycle of this parasite. Dogs and cats may also become infected if they eat rodents carrying the parasite. When eggs of *Echinococcus* are passed in the feces of the dog and cat, humans are at risk for infection. Free-roaming cats and dogs may need to be periodically treated with tapeworm medication. Rodent control and good hygiene are important in preventing the spread of this disease to humans. As with the more common tapeworm, infection with *Echinococcus* is infrequent yet possible.

*This client information sheet is based on material written by Ernest Ward, DVM
© Copyright 2005 Lifelearn Inc. Used with permission under license. May 24, 2011*